
Management of Medical Emergencies in Dental Practice VASOVAGAL SYNCOPE - CLINICAL SCENARIO

(Refer Slide Time: 00:13)



Let us see a clinical scenario of management of syncope. Syncope, is an abrupt transient loss of consciousness associated with inability to maintain postural tone.

(Refer Slide Time: 00:28)





In this video, we see the dental surgeon is in conversation with her clinic assistant and reluctant to stop the conversation even after the patient is inside the consultation cabinet. This behaviour affects the patient doctor relationship.

(Refer Slide Time: 00:47)



Dental surgeon has not taken any history from the patient and started the treatment without developing a rapport with the patient.

(Refer Slide Time: 01:05)



The patient looks quite anxious. The dental surgeon started loading the syringe with local anaesthesia in front of the patient which further led to an increase in the anxiety of the patient. This can be avoided.

(Refer Slide Time: 01:30)





The dental surgeon started the treatment without observing the patient and not even considering the patient's anxiety. When the patient had the syncopal attack, the entire team is panicking. The attending dental surgeon was unaware of the pre-syncopal symptoms and the attender running inside the clinic which should not be happening. These clinical scenarios can be avoided when you manage a patient in the dental office.

(Refer Slide Time: 02:03)





Now, let us see how to manage a patient in the dental office. The dental surgeon attended the patient immediately coming down the patient. The dental surgeon started taking a detailed case history from the patient before starting the examination. Then the dental surgeon slowly started explaining about the treatment to the patient.

(Refer Slide Time: 02:45)



The treatment was started with prior preparation. The treatment steps were explained to the patient.

(Refer Slide Time: 03:00)



Even then, the patient is going for a syncope. The dental surgeon is talking to the patient and reassuring the patient.

(Refer Slide Time: 03:10)



The attending team is very much ready to handle the situation. Clinical assistant was quick to respond in the scenario. The patient was put in supine position with legs elevated and the vitals are monitored.

(Refer Slide Time: 03:52)





The assistant was ready with a monocle spirit, which acts as a respiratory stimulant in the initial management of the patient who has gone into syncope. The patient recovered, following which the procedure was done and the patient discharged.